



ACTIVE INGREDIENTS:

Didecyl Dimethyl Ammonium Chloride.....	1.268%
n-Alkyl (C14 50%, C12 40%, C16 10%)	
dimethyl benzyl ammonium chloride.....	0.845%
INERT INGREDIENTS:	97.887%
TOTAL.....	100.000%

EPA REG. NO. 10324-157-84179 EPA EST. NO. 84179-NJ-1

disinfex lemon™

SCENT: FRESH CITRON

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield, rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse. ENVIRONMENTAL HAZARD This product is toxic to fish. PHYSICAL OR CHEMICAL HAZARDS Do not mix with oxidizers, anionic soaps and detergents. Do not use or store near heat or open flame.

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

net contents: 1 gallon



Santec Inc. 1420 E. Linden Avenue, Linden NJ 07036 P 908-912-2500
F 908374-7149 Emergency # 800-424-9300 W:www.santecchemical.com

- This product is a multi-purpose, neutral pH, germicidal detergent and deodorant effective in hard water up to 400 ppm. Disinfects, cleans, and deodorizes in one labor-saving step. For use in Hospitals, Nursing Homes, Hotels, and Schools. This product can be used on hard, non-porous surfaces such as: countertops, floors, walls, glazed tile, glazed porcelain and other hard non-porous surfaces.
- **DIRECTIONS FOR USE:**
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- Before using this product, food products and packaging materials must be removed from the room or carefully protected.
- **GENERAL DEODORIZATION:** To deodorize, add 4 ounces to one gallon of water. Allow to air dry. **DISINFECTION PERFORMANCE IN INSTITUTIONS AND NON-MEDICAL FACILITIES:** At 4 fluid ounces per 1 gallon of water use level and with a 10-minute contact time, this product is effective against the following organisms on hard non-porous surfaces: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus
- **FUNGICIDAL PERFORMANCE:** When used as directed at 4 fluid ounces per 1 gallon of water and a 10-minute contact time, this product is highly effective against the following on hard non-porous surfaces: Aspergillus niger, Candida albicans, Trichophyton mentagrophytes
- **PREPARATION OF USE SOLUTION:** Mix 4 ounces with 1 gallon of water. **TOILET BOWLS/URINALS:** Swab bowl with brush to remove heavy soil prior to cleaning or disinfecting. Clean by applying use-solution around the bowl and up under the rim. Stubborn stains may require brushing.
- To disinfect, first preclean to remove heavy soil, then remove or expel the residual bowl water over the inner trap. Pour in ½ to 1 oz. of the use solution to thoroughly cover surfaces. Swab the bowl completely using a scrub brush or mop, making sure to get under rim. Let stand for 10 minutes or overnight, then flush. Refill toilet bowl or urinal with water. Prepare a fresh solution at least daily or when use solution becomes visibly soiled.
- **MOLD AND MILDEW CONTROL:** To control the growth of mold and mildew and their odors on hard, non-porous surfaces, apply use solution of 4 ounces per 1 gallon of water. Apply by cloth, sponge, mop, brush, auto scrubber, mechanical spray device or trigger spray device to thoroughly wet all treated surfaces completely. Allow surface to remain wet for ten minutes. Allow solution to dry on surface and repeat treatment every seven days or when new growth appears. Change cloth, sponge or towels frequently to avoid redeposition of soil. Prepare a fresh solution at least daily or when use solution becomes visibly soiled. This product is not for use on medical device surfaces. This product is a Hospital Disinfectant at 4 fluid ounces per gallon with a 10 minute contact time. **PREPARATION OF DISINFECTION/FUNGICIDAL/VIRUCIDAL* USE SOLUTION:** For heavily soiled areas, a preliminary cleaning is required. Add 4-ounces with 1 gallon of water to disinfect hard non-porous surfaces. Apply use solution using a brush, cloth, mop, sponge, auto scrubber, mechanical spray device, trigger spray device or by immersion so as to thoroughly wet surfaces. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or sponge.
- Change cloth, sponge or towels frequently to avoid redeposition of soil. Treated surfaces must remain wet for 10 minutes. Allow to air dry. Rinse all treated surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse.
- Prepare a fresh solution at least daily or when use solution becomes visibly soiled. *KILLS HIV, HBV AND HCV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/ objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) associated with AIDS, Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.** Personal Protection: Clean up must always be done wearing protective latex gloves, gowns, masks and eye protection. Cleaning Procedure: Blood and other body fluids containing HIV-1, HBV and HCV must be thoroughly cleaned from hard nonporous surfaces and objects before application of this product. Disposal of Infectious Materials: Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. Contact Time: Leave hard nonporous surfaces wet for 10 minutes with 4 fluid ounces per gallon use solution.
- **STORAGE AND DISPOSAL**
- Do not contaminate water, food, or feed by storage and disposal. **CONTAINER DISPOSAL/PESTICIDE STORAGE:** Non-refillable container. Store only in original container. Do not reuse empty container. Store pesticides away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use. Triple rinse as follows: Fill container ¼ full with water and reclose the container. Agitate vigorously and dispose of rinsate. Repeat two more times.
- Wrap container and put in trash or offer for recycling if available. **RESIDUE REMOVAL INSTRUCTIONS:** Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Follow Pesticide Disposal instructions for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.