

st sapphire™

santec® ^one step cleaning and disinfecting

^^One Step disinfectant* claim is limited to association with non-food contact surface use.

General Disinfectant ready-to-use spray. This product is intended for use in airports, ambulances, athletic facilities, household bathrooms, institutional bathrooms, correctional facilities, daycare centers, gyms, health clubs, healthcare institutions, homes, household kitchens***, institutional kitchens***, laundry rooms, nursing homes and schools.

- **Kills 99.9% Of Bacteria In 5 Seconds!*****
- **Kills Cold^{††} and Flu Viruses**
- **Eliminates Odors**
- **This product is bactericidal in the presence of organic soil against Methicillin Resistant Staphylococcus aureus (MRSA) at 2 minutes contact time.**

††Rhinovirus **Staphylococcus aureus (Staph), Klebsiella pneumoniae

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **IF ON SKIN:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

ACTIVE INGREDIENTS:

Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	0.15%
Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides.....	0.15%
INERT INGREDIENTS	99.70%
TOTAL	100.00%

EPA REG. NO. 1839-220-84179 EPA EST. NO. 84179-NJ-1
Sold by: Santec, Inc. 1420 East Linden Ave, Linden NJ 07036

net contents: 1 Quart (32 fl. oz.)



Kills 99.9% of Staphylococcus aureus (Staph) and Klebsiella pneumoniae in 5 seconds. ***When using on food contact surfaces: thoroughly rinse all treated surfaces with potable (tap) water.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. **DISINFECTING, DEODORIZING:** Remove gross filth or heavy soil prior to application of the product. Apply product with a low pressure coarse sprayer. Hold sprayer six to eight inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 2 minutes. No scrubbing necessary. Allow treated surfaces to air dry or wipe off with a clean cloth. When using on food contact surfaces: countertops, stove tops, highchairs, kitchen appliances, thoroughly rinse all treated surfaces with potable water. This product must not be used to clean eating utensils, glassware and dishes. This product is an effective disinfectant against: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Methicillin Resistant Staphylococcus aureus (MRSA), Vancomycin Resistant Enterococcus faecium, Multi-Drug Resistant (MDR) Acinetobacter baumannii Ceftazidime and Gentamicin Resistant, Extended Spectrum Beta-Lactamase (ESBL) Escherichia coli and Klebsiella pneumoniae New Delhi Metallo-Beta Lactamase (NDM-1) Carbapenem Resistant. **SANITIZE NON-FOOD CONTACT SURFACES AND DEODORIZE:** To sanitize/deodorize hard, non-porous surfaces, hold container six to eight inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 5 seconds. This product is an effective sanitizer against Staphylococcus aureus (Staph) and Klebsiella pneumoniae. Let surface dry. For heavily soiled surfaces, a pre-cleaning step is required. This product will not harm most surfaces; including acrylic, sealed fiberglass and vinyl. **Do not use on unpainted wood. TO DISINFECT FOOD SERVICE ESTABLISHMENT FOOD CONTACT SURFACES:** countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 2 minutes. Wipe off with a clean cloth, mop or sponge and thoroughly rinse all food contact surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

GENERAL CLEANING: Spray soiled area, then wipe with a dry paper towel or lint-free cloth. **VIRUCIDAL ACTIVITY:** This product is an effective virucide on hard, non-porous surfaces against Human Immunodeficiency Virus Type-1 (HIV-1) and Avian Influenza Virus (H5N1) NIBRG-14, when the treated surface is allowed to remain wet for 30 seconds. This product is an effective virucide on hard, non-porous surfaces against Human Rotavirus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Duck Hepatitis B Virus (DHBV) and Bovine Viral Diarrhea Virus (BVDV) when the treated surface is allowed to remain wet for 1 minute. This product is an effective virucide on hard, non-porous surfaces against Feline Calicivirus, Norovirus (Norwalk Virus), Canine Parvovirus, Rhinovirus Type 14 and Rhinovirus Type 39 when the treated surface is allowed to remain wet for 2 minutes. This product is effective against Poliovirus Type 1 when the treated surface is allowed to remain wet for 5 minutes. Allow treated surfaces to air dry, or wipe off with a clean cloth. **KILLS HIV-1, HBV and HCV ON PRE-CLEANING ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1, Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surfaces to remain wet for 2 minutes. HIV-1 is inactivated in 30 seconds; HBV and HCV are inactivated in 1 minute. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

PRECAUTIONARY STATEMENTS Hazards To Humans And Domestic Animals

Caution: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

STORAGE AND DISPOSAL DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. **PESTICIDE STORAGE:** Store in a dry place inaccessible to children. **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose in a sanitary landfill, by incineration or burning, if allowed by state and local authorities. If burned, stay out of smoke. **PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.